

G:\Data\CLIENT\2039\004\App-PreAmerd.doc

In Vitro Selection and Optional Identification of  
Polypeptides Using Solid Support Carriers

BACKGROUND OF THE INVENTION

1. Field of the Invention

This invention provides methodology for in vitro selection and, if desired, subsequent identification of proteins or peptides with desired properties from pools of protein or peptide variants (libraries).

2. Discussion of the Related Art

Proteins and peptides, hereinafter jointly referred to as polypeptides, with desired properties such as binding affinity to a particular target molecule, catalytic activity, chemical or enzymatic activity or immunogenic activity are of great importance in many areas of biotechnology such as drug and vaccine development, diagnostic applications and bioseparation.

Recent progress in gene technology has provided the introduction of novel principles of isolating and identifying such polypeptides from large collections of variants constructed by different methods including combinatorial principles (Clackson and Wells, *Trends Biotechnol.* **12**, pp. 173-184 [1994]). Typically, using biosynthesis for production of the library members, large pools of genes are constructed, encoding the individual library members, allowing for later selection or enrichment of desired variants using an appropriate bait molecule or chemical condition (Smith and Petrenko, *Chem. Rev.* **97**, pp. 391-410 [1997]). For identification of selected variants, several techniques have been described to provide a physical link between the translated protein (phenotype) and the genetic information encoding it (genotype), allowing for identification of selected library members using DNA sequencing technology.

Using phage or cell display technologies, a genotype-phenotype coupling is obtained through

incorporation of the individual library members into the coat or cell surface structures respectively of phage or cells containing the corresponding gene, which is typically inserted into phage, phagemid, plasmid or viral DNA. In the construction of such libraries, the gene pools need to be transformed into a recipient cell used for biosynthesis of the corresponding proteins. The practical limitations associated with this critical step to obtain large (complex) libraries (typically above  $10^9$  different members) have been a driving force for the development of alternative technologies based on *in vitro* transcription and translation of genetic information, thereby avoiding the transformation step.

Examples of such technologies are ribosomal display (Mattheakis *et al.*, *Proc. Natl. Acad. Sci. USA* **91**, pp. 9022-9026 [1994]; Hanes *et al.*, *FEBS Letters* **450**, pp. 105-110 [1999]) and RNA-peptide fusions using puromycin (Roberts and Szostak, *Proc. Natl. Acad. Sci. USA* **94**, pp. 12297-12302 [1997]). In ribosomal display, a gene pool (typically polymerase chain reaction (PCR) products containing signals necessary for transcription and translation) is transcribed *in vitro* to produce a corresponding pool of mRNA used for ribosome mediated translation of proteins which typically, through the absence of translational stop signals, remain physically linked to the ribosome-mRNA complex. This allows for selection of polypeptides on the basis of the characteristics of the same and identification through DNA sequencing after conversion of the ribosome-associated mRNA into DNA by the use of reverse transcriptase. However, special precautions (temperature, buffer conditions) must be taken to ensure the stability of the ribosome-mRNA-protein complexes, limiting the conditions under which selection can be performed (Jermutus *et al.*, *Curr. Opin. Biotechnol.* **9**, pp. 534-548 [1998]; Hanes *et al.*, *op. cit.* [1999]). In

the RNA-peptide fusion system, puromycin-tagged RNA is used during translation, resulting in covalent RNA-protein/peptide links via acceptance by the ribosome of puromycin in the nascent polypeptide chain. However, 5 new puromycin-mRNA fusions have to be prepared for each round of selection, severely limiting the efficiency of the technology (Jermutus et al., *op. cit.* [1998]; Roberts, *Curr. Opin. Chem. Biol.* 3, pp. 268-273 [1999]).

A further system has been described by Tawfik and 10 Griffiths (*Nature Biotechnology*, (1998) 16; 652-656) which is cell free but seeks to mimic the effect of cells in creating compartments to link genotype and phenotype. Micelles are formed using a water-in-oil emulsion which can then be broken by mixing with ether. However, this 15 system is not without problems, the two phase system results in several practical limitations. In order to recover the encapsulated molecules, the two phase system must be broken which is rather laborious, requiring several washes and causing a loss of material. 20 Furthermore, the non-water components necessary to create the two-phase system might inhibit or denature biomolecules and the encapsulation itself makes it more difficult to deliver additional reagents necessary for e.g. detection or capture of specific molecular entities.

25 OBJECTS AND SUMMARY OF THE INVENTION

The present invention is based on the finding that by using a solid support such as a particle system as carrier of genetic information (e.g. RNA or DNA) used for identification and having coupled thereto the 30 corresponding *in vitro* translated polypeptide, methodology linking genotype and phenotype is established. Isolation of solid support particles carrying a desired library member or members may typically be performed using sorting technology employing 35 e.g. fluorescent labels incorporated into a target molecule or the library polypeptide members or by

magnetic isolation using magnetic particles containing an immobilized target molecule.

Thus according to one aspect of the present invention there is provided a method for the selection of  
5 one or more desired polypeptides comprising:

- (a) cell free expression of nucleic acid molecules immobilized on a solid support system to produce polypeptides, the solid support carrying means for biospecific interaction with at least the desired  
10 polypeptide or a molecule attached thereto;
- (b) separation of the solid support carrying both the desired polypeptide and the nucleic acid encoding it; and optionally
- (c) recovery of the said nucleic acid and/or said  
15 desired polypeptide, preferably of the nucleic acid.

The selection method of the invention can be considered also as a method of enriching the desired polypeptide from a starting library of molecules containing it. 'Enrichment' referring to increasing the  
20 relative proportion of the desired polypeptide within the sample of variant molecules. Similarly, the method can be considered one by which a nucleic acid molecule of interest, i.e. which encodes the desired polypeptide is enriched.

25 Step (a) is cell free. The term "cell" is used in a broad sense to include cell and preferably cell-like systems and thus preferably encompasses liposomes, micelles formed by water-in-oil emulsions, gels, glass or any other multi-phase system which creates a physical  
30 barrier between one gene expression/biospecific interaction system and another. According to a preferred aspect of the present method, no actual compartmentalisation takes place, no membrane or other separation system is required to isolate individual  
35 nucleic acid molecules from one another.

The separation step (b) may advantageously be effected by interaction of the immobilized desired

polypeptide with a target (e.g. biospecific) reactant therefor which carries means permitting separation of the resulting solid support/nucleic acid/desired polypeptide/target reactant complex. Such means may, for example, comprise a label such as a fluorescence label or a magnetic particle. In this way the complex may be separated using fluorescence-activated cell sorting (FACS) technology or magnetic separation technology.

The immobilized nucleic acids may, for example, be RNA or DNA encoding individual polypeptides such as the members of a protein library. It will be appreciated that their *in vitro* translation will be effected in combination with or following *in vitro* transcription in the case of immobilised DNA.

Suitable solid supports for use in the present invention may be any of the well known supports or matrices which are currently widely used or proposed for immobilisation, separation etc. These may take the form of particles, sheets, gels, filters, membranes, fibres, capillaries, or microtitre strips, tubes, plates or wells etc., particulate solid supports being preferred. Conveniently the support may be made of glass, silica, latex or a polymeric material.

Non-magnetic polymer beads suitable for use in the method of the invention are available from Dyno Particles AS (Lillestrøm, Norway) as well as from Qiagen, Pharmacia and Serotec. However, to aid manipulation and separation, magnetic beads are preferred. The term "magnetic" as used herein means that the support is capable of having a magnetic moment imparted to it when placed in a magnetic field, and thus is displaceable under the action of that field.

Thus, using the method of the invention, after gene expression and biospecific interaction the magnetic particles may be removed onto a suitable surface by application of a magnetic field eg. using a permanent magnet. It is usually sufficient to apply a magnet to

the side of the vessel containing the sample mixture to aggregate the particles to the wall of the vessel and to pour away the remainder of the sample.

Especially preferred are superparamagnetic particles  
5 for example the well-known magnetic particles sold by Dynal AS (Oslo, Norway) as DYNABEADS, are suited to use in the present invention.

Methods for attachment of nucleic acid molecules or proteinaceous moieties such as the cognate binding  
10 partners or target molecules discussed herein to a solid support are well known in the art and many include but are not limited to chemical coupling, e.g. involving amine, aldehyde, thiol, thioether or carboxyl groups or biospecific coupling for example taking advantage of  
15 interactions between streptavidin and biotin or analogues thereof, IgG and protein A or G, HSA and protein G, glutathione S-transferase (G-ST) and glutathione, maltose and maltose binding protein, antibody and antigen  
20 (including proteins, peptides, carbohydrates and haptens), lectins and carbohydrates, hisidines and chelating groups and nucleic acid/nucleic acid hybridization.

The expressed polypeptides may advantageously be fusion proteins containing an affinity fusion partner,  
25 the solid support carrying a cognate binding partner for said affinity fusion partner as the means for biospecific interaction. Thus the expressed fusion protein will typically comprise an affinity fusion partner portion as well as the desired polypeptide or a molecular variant of  
30 the desired polypeptide from the library of molecules which contains the desired polypeptide. In this way a library of fusions proteins is generated having a variable portion which is made up of the desired polypeptide or variants thereof from the starting library  
35 and an essentially common portion, the affinity fusion partner. As appropriate, reference is made herein to molecular libraries which may be libraries of nucleic

- 7 -

acid molecules or libraries of polypeptides. Likewise, a library member may refer to a polypeptide or a nucleic acid molecule.

5 In an alternative embodiment a target molecule  
capable of biospecific interaction with the desired  
polypeptide is immobilized on the solid support as the  
means for biospecific interaction. In this embodiment, a  
library of fusion proteins may also be generated, each  
fusion protein incorporating a reporter protein which may  
10 conveniently be used in the separation step (b) as well  
as the desired polypeptide or a molecular variant of the  
desired polypeptide from the library of molecules which  
contains the desired polypeptide. Thus again, the motif  
of a variable portion and an essentially common portion  
15 (here the reporter protein) is provided. Each molecule  
within the library of fusion proteins will thus  
preferably have a region which is essentially the same as  
the corresponding region of other molecules in the  
library, while the variable region of each library member  
20 will differ from all or at least most of the  
corresponding regions of the other library members. In  
general one variable region will not differ significantly  
from some or all of the other variable regions within the  
library of fusion proteins. In this way the impact of  
25 minor variations in primary amino acid sequence on e.g.  
binding can be investigated.

Recovery of the nucleic acid(s) encoding the desired  
polypeptide(s) may, for example, be effected by *in vitro*  
amplification, e.g. by means of PCR, reverse  
30 transcriptase PCR or rolling circle amplification.

The sequence of separated and/or amplified nucleic  
acid(s) may be determined, e.g. by conventional  
sequencing techniques, thereby permitting determination  
of the sequence of the desired polypeptide in order to  
35 identify it.

In a further aspect of the invention the starting  
pool of nucleic acids encoding individual library members

Performance of repeated cycles in accordance with the invention, optionally employing solid phase support particles with successively decreasing numbers of nucleic acid anchoring sites, and optionally with simultaneous dilution of the nucleic acid material, may result in gradual convergence to a limited set of library members which may be subjected to individual analysis at a clonal level in order to identify a desired polypeptide species.

Alternatively, after a reduction in the number of library members by separation in accordance with the invention, the enriched pool of nucleic acid sequences may be subjected to further selections using a different selection principle, such as (but not limited to) cell display, phage display, plasmid display, ribosomal display or mRNA-peptide fusions.

Thus, the method of the invention is preferably an iterative process with enrichment of the polypeptide(s) of interest occurring as more cycles are performed. While there may be some diffusion of expressed polypeptides and binding to neighbouring beads (or



regions of solid support, particles etc.), local binding to the polypeptide's own bead (or region of solid support, particle etc.) will be preferred. Thus after several cycles significant enrichment will be achieved.

5 Method steps (a) and (b) will thus preferably be performed more than once, typically the number of cycles will be between 1 and 100, preferably 2 to 50, more preferably 2 to 20, e.g. 5 to 10. In this way the number of variants may be very significantly limited and the  
10 relatively small number remaining can be analysed one-by-one, e.g. by ELISA, statistical analysis of clones after sequencing or Biacore analysis.

In another aspect of the invention, the selection of a solid phase support carrier carrying multiple nucleic  
15 acid species, including the desired library member, may be used to produce useful reagents without the need for identification of the particular desired library member.

Thus the method may be performed in an iterative manner but stopped when the selected sample still contains a  
20 mixed population of DNA molecules; this pool of DNA fragments can be used as a "polyclonal" material, not defined at the molecular level but still useful.

In a further embodiment of the invention two different nucleic acid libraries may be immobilized on  
25 separate solid support systems and the method may be used to select and identify interacting pairs of polypeptides.

Thus, for example, one of the nucleic acid libraries may encode polypeptides such as antibodies, antibody  
fragments, peptides or protein domains and the other may  
30 encode cDNA encoded polypeptides.

According to a further aspect of the invention is provided a molecular library comprising a solid support system having immobilised thereon a plurality of nucleic  
acid molecules and associated with each of said nucleic  
35 acid molecules and also immobilised on said support system means for biospecific interaction with the expression product of one or more of said nucleic acid molecules.

- 10 -

The solid support system is preferably particulate and thus each particle will conveniently carry one nucleic acid molecule from the library and means for biospecific interaction with the expression product  
5 thereof. Thus the aforementioned 'association' between nucleic acid molecules and means for biospecific interaction is achieved. As discussed in more detail above, the library of nucleic acid molecules will conveniently encode fusion proteins and the means for  
10 biospecific interaction may interact, typically bind, to either the variable or common portion of said fusion protein.

#### BRIEF DESCRIPTION OF THE DRAWINGS

In the accompanying drawings, which serve to  
15 illustrate the invention without in any way limiting it:

**Fig. 1** is a schematic description of the basic concept of the invention. A pool of nucleic acid fragments encoding individual polypeptide library members are immobilized onto particles of a solid support  
20 carrier. In a DNA-based format, fragments are immobilized whereafter a coupled transcription/translation step is performed resulting in the production of the corresponding gene products. In an RNA-based format, RNA molecules are transcriptionally produced from  
25 the DNA fragments, after which they are immobilized onto the solid support carrier, followed by a translation step resulting in the corresponding gene products. Typically, but not exclusively, the gene products are fusion proteins between polypeptide library members and an  
30 affinity fusion partner for which a cognate binding partner is present on the solid support carrier particles. Functional selection of a desired polypeptide results in isolation of particles carrying the corresponding genes (DNA or RNA) which are identified  
35 after nucleic amplification and DNA sequencing.

**Fig 2** is a schematic description of the use of a solid support as carrier of coupled genetic and protein

- 11 -

information (immobilized DNA/labelled target in solution version). A library of DNA constructs (typically but not exclusively PCR fragments) containing signals necessary for library member RNA transcription and protein

5 translation is immobilized onto particles of a suitable carrier support (e.g. using biotin/streptavidin chemistry by incorporation of a biotin group into the DNA of the primer used for the PCR amplification and the use of streptavidin coated beads). The genetic constructs

10 encode individual library members as genetically fused to a common affinity fusion partner (AFP) for which the cognate binding partner (CBP) is immobilized onto the particles (e.g. via suitable coupling chemistry such as streptavidin/biotin chemistry). After addition of

15 components for *in vitro* transcription and translation (e.g. an *Escherichia coli* S30 extract), RNA (mRNA) molecules are produced which encode for the different subsequently translated protein library members. Through interaction between the immobilized binding partner and

20 the newly translated affinity fusion partner, the individual library members are physically linked to the solid support carrier particles containing the genetic information (DNA) encoding them.

After washing, the solid support carrier particles

25 are incubated with labelled target molecules, e.g. comprising fluorescein isothiocyanate (FITC), allowing physical isolation of fluorescent-positive particles, for example by FACS or by magnetic separation. Thus, particles carrying complexes between the labelled target

30 and the particle-associated library member gene product and its genetic information (DNA) are isolated.

Using e.g. PCR, the DNA fragments coupled to individual or multiple isolated particles or beads are re-amplified and used for identification of the selected

35 polypeptide(s) or optionally consecutive rounds of particle immobilization, *in vitro* transcription and translation followed by selection, e.g. by FACS.

**Fig. 3** is a schematic representation of the use of a solid support as carrier of coupled genetic and protein information (immobilized mRNA/labelled target in solution version). From a library of genetic constructs containing signals necessary for library member transcription and protein translation, RNA (mRNA) is produced (transcription) *in vitro* and immobilized onto particles of a suitable carrier support (e.g. via hybridization between complementary sequences present in the mRNA and immobilized DNA, PNA or RNA fragments). The immobilized mRNA molecules encode individual library members as genetically fused to a common affinity fusion partner (AFP) for which the cognate binding partner (CBP) is immobilized onto the particles (e.g. via streptavidin/biotin chemistry). After addition of components for *in vitro* translation (e.g. an *Escherichia coli* S30 extract), the mRNA molecules are translated to produce the different protein library members. Through interaction between the immobilized binding partner and the newly translated affinity fusion partner, the individual library members are physically linked to the solid support carrier particles containing the genetic information (mRNA) encoding them.

After washing, the solid support carrier particles  
25 are incubated with labelled target molecules, e.g.  
comprising FITC, allowing physical isolation of  
fluorescent-positive particles, e.g. by FACS. Thus,  
individual or multiple particles carrying complexes  
between the labelled target and the particle-associated  
30 library member gene product and its genetic information  
(mRNA) are isolated.

Using e.g. reverse transcriptase PCR, the bead/particle-associated mRNA molecules are converted into the corresponding DNA fragments which are PCR amplified and used for identification of the selected polypeptide(s) or optionally consecutive rounds of *in vitro* transcription, particle immobilization, *in vitro*

translation followed by selection, e.g. by FACS or magnetic selection.

Fig. 4 is a schematic representation of the use of a solid support as carrier of coupled genetic and protein information (immobilized DNA/labelled binder version). A library of DNA construct (typically but not exclusively PCR fragments) containing signals necessary for library member RNA transcription and protein translation is immobilized onto discrete particles of a suitable carrier support (e.g. using biotin/streptavidin chemistry by incorporation of a biotin group into the DNA of the primer used for the PCR amplification and the use of streptavidin coated beads). The particles also carry the target molecule with which interacting library members are desired to interact. This immobilization can be achieved using e.g. standard coupling chemistries such as EDC/NHS chemistry or biotin/streptavidin chemistry. The genetic constructs encode individual library members as genetically fused to a reporter fusion partner (RFP) such as an enzyme or autofluorescent protein such as green fluorescent protein (GFP). After addition of components for *in vitro* transcription and translation (e.g. an *Escherichia coli* S30 extract), RNA (mRNA) molecules are produced which encode for the subsequently translated different protein library members. Through interaction between the immobilized target molecule and the newly translated library member, individual library members capable of interaction with the solid support immobilized target molecule are physically linked to the solid support carrier particles containing the genetic information (DNA) encoding them.

After washing, the solid support carrier particles are sorted, e.g. using FACS technology or magnetic separation, to isolate individual or multiple particles carrying complexes between the immobilized labelled target and the particle-associated library member gene product. Thus, particles carrying complexes between the

Using PCR, the DNA fragments coupled to discrete  
5 isolated beads are re-amplified and used for  
identification of the selected polypeptide(s) or  
optionally consecutive rounds of particle immobilization,  
in vitro transcription and translation followed by  
separation, e.g. by FACS or magnetic selection.

**Fig. 5** is a schematic representation of the use of a solid support as carrier of coupled genetic and protein information (immobilized mRNA/labelled binder version). From a library of genetic constructs containing signals necessary for library member transcription and protein translation, RNA (mRNA) is produced (transcription) *in vitro* and immobilized onto particles of a suitable carrier support (e.g. via hybridization between complementary sequences present in the mRNA and immobilized DNA, PNA or RNA fragments). The particles also carry the target molecule with which library members are desired to interact. This immobilization may be obtained using e.g. standard coupling chemistries such as EDC/NHS chemistry or biotin/streptavidin chemistry. The genetic constructs (mRNA) encode individual library members as genetically fused to a reporter fusion partner (RFP) such as an enzyme or autofluorescent protein such as green fluorescent protein (GFP). After addition of components for *in vitro* translation (e.g. an *Escherichia coli* S30 extract), mRNA molecules are translated to produce the different protein library members. Through interaction between the immobilized target molecule and the newly translated library member, individual library members capable of interaction with the solid support immobilized target molecule are physically linked to the solid support carrier containing the genetic information (mRNA) encoding them.

- 15 -

After washing, the solid support carriers are sorted, e.g. using FACS technology, to isolate individual or multiple particles carrying complexes between the immobilized labelled target and the particle-associated library member gene product. Thus, particles carrying  
5 complexes between the labelled target and the particle-associated library member gene product and its genetic information (mRNA) are isolated.

Using e.g. reverse transcriptase PCR, the  
10 bead/particle-associated mRNA molecules are converted into the corresponding DNA fragments which are PCR amplified and used for consecutive rounds of *in vitro* transcription, particle immobilization, *in vitro* translation followed by selection, e.g. by FACS.

15 **Fig. 6** illustrates the experimental set-up for Example 1. Paramagnetic particles coated with streptavidin were firstly incubated with biotinylated human serum albumin (HSA), resulting in robust anchoring of HSA. Separate aliquots were subsequently incubated  
20 with either (A) protein ABD-Z, a genetic fusion protein between a serum albumin binding protein (ABD) derived from streptococcal protein G and an immunoglobulin binding protein (Z) derived from staphylococcal protein A, followed by incubation with fluorescent isothiocyanate  
25 (FITC) conjugated polyclonal goat IgG antibodies, or (B) with the FITC conjugated goat IgG antibodies directly.

**Fig. 7** is a photograph from UV-microscopy analyses of streptavidin-coated beads/particles containing streptavidin/biotin chemistry-immobilized biotinylated  
30 human serum albumin. (A) Particles incubated with FITC-conjugated polyclonal goat IgG antibodies after having first been subjected to a solution containing the fusion protein Z-ABD. (B) Particles incubated with FITC-conjugated polyclonal goat IgG antibodies only.

35 **Fig. 8** is a schematic representation of the use of the invention for selecting interacting polypeptide pairs through the crossing of two different libraries. Two

pools of nucleic acid fragments encoding different polypeptide libraries are separately immobilized onto particles of solid support carrier systems. In a DNA-based format, fragments are immobilized whereafter a coupled transcription/translation step is performed resulting in the production of the corresponding gene products. In an RNA-based format, RNA molecules are transcriptionally produced from the DNA fragments, after which they are immobilized onto the solid support carrier, followed by a translation step resulting in the corresponding gene products. Typically, but not exclusively, the gene products are fusion proteins between polypeptide library members and an affinity fusion partner for which a cognate binding partner is present on the particles. The different libraries are differently labelled, e.g. using two fluorophores having different excitation spectra. Biospecific interactions between members of the different polypeptide libraries are detected as double-labelled particle pairs. For identification, the nucleic acids present on the isolated particles encoding the corresponding genes are analyzed by DNA sequencing.

**Fig. 9** is a schematic description of the construction of the plasmids pGEM-SD-K-FLAG-Z<sub>wt</sub> and pGEM-SD-K-FLAG-Z<sub>IgA</sub>, designed for use as template for the amplification of PCR products for cell free transcription and translation of either free or bead-immobilized DNA/RNA.

**Fig. 10** is a radiograph obtained after SDS-PAGE analysis under reducing conditions of proteins synthesized using a cell free extract supplemented with [35S]methionine and PCR products produced with primers NOOL-12 and NOOL-13 using different plasmids as templates. Lane 1: pGEM-SD-K-FLAG-Z<sub>wt</sub>, lane 2: pGEM-SD-K-FLAG-Z<sub>IgA</sub>. A marker with 14C-labeled proteins was used as size reference (prod. no. CFA756, Amersham Pharmacia Biotech, Uppsala, Sweden). Arrows indicate the



- 17 -

positions of reference proteins with molecular weights of 14.3, 20.1 and 30.0 kDa, respectively.

**Fig. 11** is an overlay plot from a comparative FACS analysis of anti-FLAG BioM5 antibody-coated beads  
5 subjected to a FLAG-Z<sub>wt</sub> PCR product transcription/translation mixture and negative control beads treated in the same way but not coated with anti-FLAG BioM5 antibodies.

**Fig. 12** is an overlay plot from a comparative FACS  
10 analysis of anti-FLAG BioM5 antibody and PCR product doubly coated beads, subjected to a transcription/translation mixture, followed by detection. The picture shows the analysis of two different sets of beads containing either FLAG-Z<sub>wt</sub> or FLAG-Z<sub>IgA</sub> encoding PCR  
15 products subjected to the analysis.

**Fig. 13 (A)** is a schematic representation of the presence of a Mlu I restriction site in the PCR product obtained by PCR amplification using primers NOOL-12 and NOOL-13 on a pGEM-SD-K-FLAG-Z<sub>wt</sub> plasmid template. In  
20 contrast, no Mlu I site is present in the PCR product obtained by PCR amplification using primers NOOL-12 and NOOL-13 on a pGEM-SD-K-FLAG-Z<sub>IgA</sub> plasmid template. Also shown are the sizes of the cleavage products obtained after incubation of the FLAG-Z<sub>wt</sub> fusion protein encoding  
25 PCR product after incubation with Mlu I.

(B) are photographs showing agarose gel electrophoresis analyses of PCR products obtained by PCR amplification of different samples taken before or after FACS-based enrichments. Lane 1: Beads containing FLAG-Z<sub>wt</sub>  
30 encoding PCR product only; lane 2: Beads containing FLAG-Z<sub>wt</sub> encoding PCR product only. Resulting PCR product subjected to incubation with Mlu I; lane 3: Beads containing FLAG-Z<sub>IgA</sub> encoding PCR product only; lane 4: Beads containing FLAG-Z<sub>IgA</sub> encoding PCR product only.  
35 Resulting PCR product subjected to incubation with Mlu I; lane 5: Beads containing a 1:1 mixture of FLAG-Z<sub>wt</sub> and FLAG-Z<sub>IgA</sub> encoding PCR products. Sample from before FACS

- 18 -

enrichment experiment; lane 6: Beads containing a 1:1 mixture of FLAG-Z<sub>wt</sub> and FLAG-Z<sub>IgA</sub> encoding PCR products. Resulting PCR product subjected to incubation with Mlu I.

Sample from before FACS enrichment experiment; lane 7:  
 5 Sample from beads sorted in FACS enrichment experiment; lane 8: Sample from beads sorted in FACS enrichment experiment. Resulting PCR product subjected to incubation with Mlu I. Flanking lanes with size markers (phage 1 DNA cleaved with Pst I, Amersham Pharmacia  
 10 Biotech, Uppsala, Sweden) are labeled M.

**Fig. 14** Top: is an overlay plot of intensity recordings of tracks corresponding to lanes 6 (dashed line) and 8 (solid line) in figure 13. The relative intensity is shown as a function of the migration  
 15 coordinate. Bottom: shows digitally excised tracks from the gel image corresponding to lane 6 and 8 from the gel shown in figure 13. A relative shift of intensity towards the smaller molecular weight cleavage products is observed for the sample obtained by PCR amplification of  
 20 nucleic acids present on beads collected in the FACS enrichment (track corresponding to lane 8).

#### DESCRIPTION OF THE PREFERRED EMBODIMENTS

In a representative embodiment of the method of the invention a pool of gene fragments (Figs. 2-5) containing  
 25 the DNA encoding different polypeptide library members is prepared using standard DNA technology, for example as described by Nord et al., *Prot. Engineering* **8**, pp. 601-608 [1995] and Nord et al., *Nature Biotechnol.* **15**, pp. 772-777 [1997]. The gene fragments should include a  
 30 first sequence corresponding to a suitable RNA polymerase promoter sequence, such as *E. coli* phage T7 promoter, T3 promoter, SP6 promoter, *lac* promoter, *lac* UV5 promoter, *ara* B promoter, *trp* promoter, staphylococcal protein A promoter, or viral promoters such as Raus Sarcoma Virus  
 35 (RSV) promoter, and Cytomegalo virus (CMV) late and early promoters to function as signals for transcription of the DNA fragment into mRNA using a suitable extract such as

- 19 -

an S30 extract of *E. coli* for promoters of *E. coli* or prokaryotic origin or a reticulocyte extract or wheat germ extract for promoters of eukaryotic origin (coupled systems) or by a first transcriptional step using a  
5 preparation of purified suitable RNA polymerase, separated from a later translational step (uncoupled system) in which the mRNA templates are used for translation of the genetic information into the corresponding polypeptides.

10 In one aspect of the invention, the promoter sequence is followed by a sequence encoding an affinity fusion partner (AFP), employed for binding a cognate binding partner immobilized onto a solid phase carrier particle. This affinity fusion partner may for example  
15 be the albumin binding region of streptococcal protein G or derivatives thereof, the immunoglobulin binding protein A or derivatives thereof, maltose binding protein, glutathione S-transferase, FLAG peptide, Bio-tag (biotinylated peptide), hexahistidyl sequence, c-myc tag,  
20 or any other polypeptide for which a suitable cognate binding partner is available. The gene fragments should each also contain the gene encoding an individual library member polypeptide, in translational frame with the affinity fusion partner polypeptide if used.  
25 Alternatively, the gene encoding the affinity fusion partner may be positioned after the gene for the polypeptide library member.

In one aspect of the invention, the sequence encoding the individual library member polypeptide is  
30 either preceded or followed by a sequence encoding a suitable reporter polypeptide, such as green fluorescent protein (GFP), alkaline phosphatase, luciferase, horse radish peroxidase (HRP) or  $\beta$ -galactosidase.

In one aspect of the invention, the gene fragments  
35 contain a suitable chemical group (e.g. biotin or digoxin) introduced e.g. by PCR amplification using a primer or nucleotides labelled with the group. This

- 20 -

group is used for anchoring the DNA fragment onto solid support particles coated with a suitable cognate binding partner, such as streptavidin or anti-digoxin antibody(ies) (Figs. 2 and 4).

5 In another aspect of the invention, a pool of transcribed mRNA is immobilized onto the solid support particles via a suitable attachment moiety. This moiety may for example be a nucleotide sequence at the 5'- or 3'- end of the mRNA, for which a complementary sequence  
10 of RNA, DNA or PNA is immobilized onto the solid support particles (Figs. 3 and 5).

After immobilization of DNA fragments onto the solid support particles, a transcription step is performed using a suitable RNA polymerase depending on the promoter  
15 used for the construction of the fragments. The thereby transcribed mRNA is employed for translation of the genetic information into the corresponding polypeptides which are bound to the solid support particles by biospecific interaction with either an immobilized  
20 cognate binding partner for an affinity fusion partner encoded in translational frame with the polypeptide or via recognition of a target molecule immobilized onto the particle. For the translation a suitable extract or pure components may be used such as an E. coli S30 extract, a  
25 rabbit reticulocyte extract or a reconstituted mixture of purified essential components of a translation machinery.

Suitable particles may for example be made of polystyrene or any other polymer or mixtures of polymers, cellulose, hydroxyapatite, sepharose, dextran or silica.  
30

After immobilization of mRNA molecules onto solid support particles, the translation of these into the corresponding proteins is performed as described above. The thereby produced polypeptides are bound to the solid  
35 support particles by biospecific interaction with either an immobilized cognate binding partner for an affinity fusion partner encoded in translational frame with the

polypeptide or via recognition of a target molecule immobilized onto the particle.

To circumvent cross-over reactions, i.e. the binding of a translated polypeptide fusion protein molecule to a cognate binding partner or target molecule present on a solid support particle not also carrying the genetic information (DNA or RNA) encoding the polypeptide, the mixture may be diluted so as to prevent close proximity between particles.

Selection of particles containing a desired polypeptide or group of polypeptides may be performed by direct isolation, for example in an FACS scanner if the target is labelled with a fluorophore or if the polypeptide is genetically fused to a fluorescent protein such as green fluorescent protein. A different selection method is to use magnetic principles, using magnetic (or paramagnetic) particles coated with the target molecule of interest (Figs. 1 and 2). Alternatively, particles labelled via a specific interaction between a library member polypeptide gene product may be physically isolated using e.g. a UV- microscope.

Selection may be performed on the basis of functional properties of the encoded polypeptides, such as binding to a desired target (antibodies or other proteins or peptides, carbohydrates, organic molecules, cells, viruses, plants etc.), catalytic activity, or through proteolytic or chemical stability under certain chemical conditions.

After isolation of particles carrying a polypeptide with the desired characteristics, the nucleic acid information (DNA or RNA) present on the same particles is amplified (if necessary) by *in vitro* nucleic acid amplification methods such as reverse transcriptase PCR (if RNA), PCR (if DNA), or rolling circle replication.

If necessary, the procedure may be repeated for additional cycles of direct DNA immobilization or RNA immobilization after *in vitro* transcription of

re-amplified particle-bound nucleic acids. If further variation is desired for the next round of selection, the amplification conditions or polymerase(s) may be chosen to introduce mutations into the next pool of DNA

5 fragments.

In yet another aspect of the invention two different libraries of polypeptides are investigated for interacting pairs (Fig. 8). Particles corresponding to a library of e.g. cDNA encoded polypeptides are mixed with  
10 particles carrying members of a polypeptide library of, for example, cDNA encoded proteins, antibodies or fragments thereof, peptides or protein domains. The particles used for the immobilization of the nucleic acids are prepared such that they contain two different  
15 labels, one for each library. Isolation of interacting pairs of polypeptides resulting from biospecific interactions are isolated by e.g. FACS technology, employing detection of double-labelled particle pairs.

The method of the invention has several advantages  
20 over existing selection systems using an *in vivo* polypeptide biosynthesis step, since there is no need for transformation of the genetic material into a recipient cell. The only limitation with respect to library size (complexity) is the binding capacity of the solid support  
25 system. Furthermore, the present *in vitro* selection system uses a robust solid support as the linkage between genotype and phenotype, enabling harsh conditions to be used when selecting ligands with high affinity towards a given target molecule. As a consequence of the nucleic  
30 acids being directly immobilized on the solid support they may easily be recovered; thus, for example, if the solid support comprises magnetic beads these may be removed from the transcription/translation mixture with a magnet, thus lowering the risk of contamination with  
35 non-immobilized nucleic acids.

The following non-limitative Example serves to illustrate the invention.

**STANDARD PROCEDURES:****Cloning and PCR amplifications:**

Standard cloning work including plasmid  
 5 preparations, restriction enzyme cleavage and ligations  
 etc. was performed as described in (Sambrook, J.,  
 Fritsch, E. F. and Maniatis, T. Molecular cloning: a  
 laboratory manual, 2nd edn., Cold Spring Harbor  
 Laboratory, New York, 1989) and according to suppliers  
 10 recommendations. Restriction enzymes and ligase were  
 purchased from either MBI Fermentas, Vilnius, Lithuania  
 or New England Biolabs, MA, USA) PCR amplifications using  
 plasmids or bead-immobilized PCR products as templates  
 were performed in a GeneAmp® PCR system 9700 (PE  
 15 Biosystems, Foster City, CA, USA), using standard  
 conditions. As primers, oligonucleotides from Table 1  
 were used as specified in the examples. Typically, 5  
 pmoles of primers were used in a 30-cycle PCR  
 amplification using a buffer consisting of 0.2 mM  
 20 deoxyribonucleoside triphosphates (dNTPs), 50 mM KCl, 2  
 mM MgCl<sub>2</sub>, 10 mM Tris-HCl (pH 8.5), 0.1% Tween 20 and 0.1  
 units of AmpliTaq® DNA polymerase (PE Biosystems). A  
 standard PCR cycle had the following settings: 15 s 94°C,  
 20 s 55°C, 1 min 72°C. Standard agarose gel  
 25 electrophoresis analyses of nucleic acids were performed  
 using ethidium bromide for staining. E. coli cells used  
 for cloning and plasmid preparations were RR1DM15  
 (Rüther, U. Nucl. Acids Res. 10: 5765-5772, 1982).

30 **Table 1. List of oligonucleotide primers.**

Name	Sequence 5'-3'
NOOL-6	GGGGGGAAGCTTGGGGGGGCGCATGGCTTTAGCTGAAGCTAAAGTCTTAG
NOOL-7	CTTTGTTGAATTTGTTGTCTACGCTCGAGCTAGGTAATGCAGCTAAAATTTTCAT
NOOL-8	ATGAAATTTTAGCTGCATTACCTAGCTCGAGCGTAGACAACAAATTCAACAAAG

NOOL-9	GGGGGAATTCTTATTATTTTCGGCGCCTGAGCATCAT
NOOL-10	GGGGGAAGCTTGGGGG
NOOL-11	GGGGGAATTCTTATTATTTTCG
NOOL-12	GTTGTGTGGAATTGTGAG
NOOL-13	Biotin-AAGTTGGGTAACGCCAGG
SD KOZAK-1	AGCTTAATAATTTTGTTTAACTTTAAGAAGGAGATATAGC
SD KOZAK-2	CATGGCTATATCTCCTTCTTAAAGTTAAACAAAATTATTA
FLAG-1	CATGGACTACAAAGATGACGATGATAAAAGC
FLAG-2	TCGAGCTTTTATCATCGTCATCTTTGTAGTC

**Recombinant protein production:**

E. coli cells used for expression were either RR1DM15  
 5 (Rüther, U. Nucl. Acids Res. 10: 5765-5772, 1982) or  
 BL21DE3 (Novagen, Madison, WI, USA). Osmotic shock  
 procedures were performed as described earlier (Nygren et  
 al., J. Mol. Recognit. 1:69-74, 1988). Affinity  
 chromatography purifications of proteins on HSA and  
 10 IgG-Sepharose resins were performed as described earlier  
 (Nygren et al., J. Mol. Recognit. 1:69-74, 1988). Human  
 polyclonal IgG was supplied by Pharmacia and Upjohn AB,  
 Stockholm.

**Protein biotinylation:**

15 Human serum albumin (HSA) (prod no. A-8763, Sigma) was  
 biotinylated using EZ-Link™ Sulfo-NHS-LC-Biotin kit  
 (prod no. 21335, Pierce Chemical Company, Rodeford, IL,  
 USA).

**20 Cell free transcription and translation of PCR fragments:**

PCR products as indicated were subjected to cell free  
 transcription and translation using a commercial E. coli  
 S30 extract system for linear DNA (prod no. L1030,



- 25 -

5 Promega, Madison, WI, USA) according to the instructions by the manufacturer. For coupled transcription/translation of free (non-immobilized) PCR products, typically, 10-70 ng of PCR product was mixed with 50 µl of cell extract and incubated for 1 h at 25°C. In other experiments, PCR products were immobilized onto streptavidin coated microbeads (M280-SA, Dynal, Norway or Bang Laboratories, prod. no. CP01N/004109, where indicated). Such beads had previously been incubated with a 1.89 mg/ml solution of biotinylated BioM5 antibody (prod no. F-2922, Sigma, Saint Louis, MO, USA) directed to a FLAG peptide for affinity capture of FLAG peptide-tagged proteins. Typically, 10 ng of PCR product were mixed with 1 mg of BioM5-containing beads, which were subsequently washed two times before a coupled transcription/translation reaction was performed using 25 µl of E. coli extract.

**Protein gel electrophoresis:**

20 Sodium dodecylsulphate polyacrylamide gel electrophoresis of proteins (SDS-PAGE) under reducing conditions was performed using the Phast system (Amersham Pharmacia Biotech, Uppsala, Sweden) or in a Novex Xcell II (San Diego, CA, USA), as described by the respective suppliers.

25 **DNA sequencing:**

DNA sequencing was performed by cycle sequencing (Carothers et al., BioTechniques 7:494-499, 1989; Savolainen, P., et al., Mol. Biol. Evol. 17:474-488, 2000) using ThermoSequenase DNA polymerase (Amersham Pharmacia Biotech) and primers as indicated. Sequencing reactions were loaded onto a ABI Prism 377XL instrument (PE Biosystems, Foster City, CA, USA).

**Fluorescence-activated cell sorting (FACS) experiments:**

35 FACS analyses were performed with either a FACSCalibur, FACScan or a FACS Vantage SE instrument (Becton Dickinson, Oxnard, USA).

Where indicated, horseradish peroxidase-conjugated antibodies were used for signal amplifications, using a fluorescein tyramide reagent (Boehringer Mannheim, Germany) as described by Anton and coworkers (Anton et al., J. Histochem. Cytochem. 46:771-777, 1998).

### Example 1

#### Discrimination between solid support particles labelled with fluorescent proteins through a biospecific interaction and control solid support particles

Approximately 2 mg of streptavidin coated particles (M280-SÅ, Dynal, Norway) were incubated with 30 µl of a 2 mg/ml solution in PBS buffer (0.15 M NaCl, 20 mM phosphate, pH 7.2) of human serum albumin (HSA) (Sigma art. No. A-8763) biotinylated using a protein biotinylation kit (Pierce art. No. 21335) according to the manufacturers instructions. Particles were then either directly incubated with polyclonal goat IgG antibodies, labelled with FITC (Sigma art. No. F-9887) or first incubated with 30 µl of a 2 mg/ml solution in PBS of a fusion protein (Z-ABD) between a serum albumin binding protein (ABD) derived from streptococcal protein G and a immunoglobulin binding protein (Z) derived from staphylococcal protein A produced and HSA-affinity purified as previously described (Nord et al., op. cit. [1995], and [1997]). Between each incubation multiple (5-10) washings with PBS were performed to remove non-specifically bound proteins.

To investigate whether discrimination was possible between particles labelled by the FITC-labelled goat polyclonal antibodies via a biospecific interaction to the Z moiety of the Z-ABD fusion protein and particles not incubated with the Z-ABD fusion protein and thus incapable of binding the goat antibody, particles were

analysed by UV-microscopy using a Olympus BH2-RFCA  
microscopy at an excitation wavelength of 495 nm. The  
results shown in Fig. 5 show that a clear difference in  
fluorescent intensity can be seen between the two  
5 differently treated pools of particles (Fig. 5A and 5B).

This shows that the result of a biospecific interaction  
between an (ABD-HSA)-immobilized fusion protein and a  
labelled target protein added in solution can be  
observed.

10

### Example 2

#### Assembly and cloning of genetic constructs for cell free transcription and translation experiments

To be able to obtain PCR products encoding relevant  
15 proteins or protein library members and suitable for cell  
free transcription and translation experiments using  
solid supports as carriers for both nucleic acids and  
their corresponding encoded proteins, a genetic construct  
was assembled in the plasmid vector pGEM-4Z (Figure 9).  
20 In a splice overlap extension (SOE) PCR reaction using  
primers NOOL-10 and NOOL-11 (table 1), two gene fragments  
encoding an albumin binding protein (APB) (Larsson, et  
al., Prot. Expr. Purif. 7:447-457, 1996) and the Z domain  
(Z<sub>wt</sub>) (Nilsson et al., Prot. Engineering 1:107-113, 1987),  
25 respectively, were joined. The two fragments had  
previously been produced by separate PCR reactions using  
pT7-ABPc (ABP) (Larsson, et al., Prot. Expr. Purif.  
7:447-457, 1996) (primers NOOL-6 and NOOL-7, table 1) or  
pKN1-Z<sub>wt</sub> (Nord et al., Prot. Engineering, 8:601-608, 1995)  
30 (primers NOOL-8 and NOOL-9, table 1) as plasmid  
templates, respectively. In the SOE reaction, two  
fragments were joined resulting in an ABP-(Ser)3-Z<sub>wt</sub>  
encoding gene fragment comprising in the 5'-end  
recognition sites for the two enzymes Hin dIII and Nco I,  
35 and in the 3'-end two translational stop codons and a  
recognition site for the restriction enzyme Eco RI  
(Figure 9). This fragment was inserted by ligation as a

- 28 -

Hin dIII-Eco RI fragment into the plasmid pGEM-4Z, cleaved with the same enzymes, resulting in the construct pGEM-ABP-Z<sub>wt</sub>.

5 A fragment was assembled by the annealing of the two  
oligonucleotides SD KOZAK-1 and SD KOZAK-2 (table 1),  
resulting in a 40 bp fragment comprising an E. coli Shine  
Dalgarno (SD) sequence (for efficient E. coli  
translation) and a Kozak sequence (to facilitate  
10 expression in cell extracts from mammalian sources),  
flanked by Hin dIII and Nco I restriction sites (Figure  
9). This fragment was inserted by ligation into  
pGEM-ABP-Z cleaved with Hin dIII and Nco I, resulting in  
the plasmid vector pGEM-SD-K-ABP-Z<sub>wt</sub>. This vector was  
15 subsequently cleaved with enzymes Nco I and Xho I,  
releasing the ABP encoding fragment. The thereby  
obtained vector fragment was ligated to a FLAG peptide  
encoding gene fragment, previously obtained by annealing  
the two oligonucleotides FLAG-1 and FLAG-2 (table 1),  
20 resulting in the vector pGEM-SD-K-FLAG-Z. This vector  
thus encodes a FLAG-Z<sub>wt</sub> fusion protein, linked by a (Ser)3  
linker (Figure 9). The vector also contains an upstream  
T7 promoter which is capable of driving the transcription  
of the FLAG-Z<sub>wt</sub> fusion protein gene by the action of T7  
25 RNA polymerase. From this vector, any suitable gene  
fragment inserted between the Xho I and Eco RI sites can  
be transcribed as an mRNA operatively linked to a SD  
sequence, a Kozak sequence and a FLAG peptide encoding  
part. In addition, using primers NOOL-12 and NOOL-13  
30 (table 1), PCR products can be obtained which are  
suitable for T7 RNA polymerase driven transcription and  
are biotinylated in their 3'-ends, suitable for  
immobilization on e.g. streptavidin coated surfaces and  
other solid supports.

35

To construct the vector denoted pGEM-SD-K-FLAG-Z<sub>IgA</sub>, in  
which the Z<sub>wt</sub> encoding gene fragment has been substituted

for a gene fragment encoding the human IgA-binding protein  $Z_{IgA}$  (Gunneriusson et al., J. Bact. 1999), a  $Z_{IgA}$  encoding gene fragment was amplified using primers NOOL-8 and NOOL-9 using a plasmid pKN1- $Z_{IgA}$  template  
5 (Gunneriusson et al., J. Bact. 1999). The resulting PCR product was cleaved with restriction enzymes Xho I and Eco RI and inserted into the vector pGEM-SD-K-FLAG- $Z_{wt}$ , previously cleaved with the same enzymes. The resulting vector pGEM-SD-K-FLAG- $Z_{IgA}$  thus encodes a FLAG- $Z_{IgA}$  fusion  
10 protein, linked by a (Ser)3 linker (Figure 9).

### Example 3

#### Cell free transcription/translation of FLAG- $Z_{wt}$ and FLAG- $Z_{IgA}$ fusion proteins from their respective PCR products.

  
15

Using the plasmid vectors pGEM-SD-K-FLAG- $Z_{wt}$  and pGEM-SD-K-FLAG- $Z_{IgA}$ , respectively, for PCR amplifications using the primers NOOL-12 and NOOL-13 (table 1), PCR products were obtained of which approx. 70 ng were  
20 subjected to a one hour cell free transcription/translation at 25°C using 50 µl of an E. coli S30 cell extract (L1030, Promega, MA, USA), supplemented with [<sup>35</sup>S]methionine and 1600 units of T7 RNA polymerase. Samples of the different transcription/translation  
25 mixtures were analyzed by 10% NuPAGE (Novex, San Diego, CA, USA) under reducing conditions through the addition of 50 mM DTT (final concentration) in the sample loading buffer (NuPAGE LDS sample buffer, Novex) followed by exposure of the gel to a film (Kodak XOMAT-AR, 18x24 cm)  
30 at -70°C over night. The development of the film revealed radioactive protein of expected sizes (~8 kDa) for both the FLAG- $Z_{wt}$  and the FLAG- $Z_{IgA}$  encoding PCR products (Figure 10). This shows that the constructed plasmid vectors pGEM-SD-K-FLAG- $Z_{wt}$  and pGEM-SD-K-FLAG- $Z_{IgA}$ , both  
35 were suitable for use as templates for the amplification of PCR products capable of directing a T7 RNA polymerase driven transcription of mRNA which could be used for cell

free translation of FLAG-Z<sub>wc</sub> and FLAG-Z<sub>IgA</sub> fusion proteins in an E. coli S30 extract.

### Example 4

5 Immobilization of FLAG-Z<sub>wt</sub> and FLAG-Z<sub>IgA</sub> fusion proteins on anti-FLAG antibody-containing beads

To investigate the functionality of the FLAG peptide moieties of the fusion proteins FLAG-Z<sub>wt</sub> and FLAG-Z<sub>IgA</sub>, reaction mixtures obtained from production of the two fusion proteins from their respective PCR products using cell free transcription/translation as described in example 3 were mixed for three hours at room temperature with streptavidin coated M-280-SA dynabeads (Dyna, Norway) (50 mg) previously incubated with 5 µl of a 1.89 mg/ml solution of biotinylated anti-FLAG BioM5 monoclonal antibodies (Sigma) in PBS (0.15 M NaCl, 20 mM phosphate, pH 7.2). In the experiment, beads which had not been incubated with the biotinylated anti-FLAG BioM5 antibody solution were also included (control).

The beads were subsequently washed with PBST (PBS with 0.1% Tween 20) and analyzed using a Beckman LS6000 SC scintillator (Beckman-Coulter, Fullerton, CA, USA), under standard conditions using scintillation buffer. The measured signals from anti-FLAG BioM5-coated beads subjected to the transcription/translation mixtures corresponding to the FLAG-Z<sub>wt</sub> and FLAG-Z<sub>IgA</sub> fusion protein, respectively, were significantly higher compared to the negative controls (Table 2). This shows that fusion proteins, here exemplified by the two fusion proteins FLAG-Z<sub>wt</sub> and FLAG-Z<sub>IgA</sub>, can be produced from their respective PCR products by cell free transcription/translation containing a functional affinity fusion partner, here exemplified by the FLAG peptide, which is suitable for immobilization of the proteins to beads containing a cognate affinity partner, here exemplified by the BioM5 anti-FLAG monoclonal antibody.

Table 2. Measured scintillation signals (accumulated under 1 min) from native streptavidin (SA) beads or streptavidin beads coated with biotinylated anti-FLAG BioM5 antibody, respectively, after mixing (and subsequent washing) with transcription/translation mixtures from different samples.

Beads	Transcription/ translation mix	Signal (cpm)
native SA beads	FLAG-Z <sub>wt</sub>	4 858
BioM5 anti-FLAG coated	FLAG-Z <sub>wt</sub>	34 966
native SA beads	FLAG-Z <sub>IgA</sub>	5 959
BioM5 anti-FLAG coated	FLAG-Z <sub>IgA</sub>	43 727

10

Example 5

Cell free transcription/translation of a FLAG-Z<sub>wt</sub> encoding PCR product, biospecific immobilization of the gene product onto beads and analysis by fluorescence-activated cell sorting (FACS)

15

Cell free transcription and translation of a PCR product obtained by PCR amplification with primers NOOL-12 and NOOL-13 (Table 1) on a pGEM-SD-K-FLAG-Z<sub>wt</sub> plasmid template was performed as in example 3, but without the addition of [<sup>35</sup>S]methionine. The resulting mixture was incubated for 2 hours with 50 mg streptavidin-coated polystyrene beads with a diameter of approximately 0.95 mm) (Bangs Laboratories, Fishers, IN, USA), previously incubated with 5 µl of a 1.89 mg/ml solution of biotinylated anti-FLAG BioM5 monoclonal antibodies. In the experiment, beads not coated with the biotinylated BioM5 anti-FLAG antibody were also included, as a control. After thorough washing with TNT buffer (0.1 M Tris-HCl pH 7.5, 0.15 M NaCl, 0.05% Tween 20), rabbit anti-DNP IgG

20

25

30

- 32 -

antibodies conjugated to horse-radish peroxidase (HRP) (art. no. P0402, Dako, Denmark) were added to the beads and incubated for 45 min at 25°C, followed by washing with TNT buffer, to detect the translated and biospecifically  
5 immobilized FLAG-Z<sub>wt</sub> fusion protein gene product via the biospecific interaction between the constant parts (Fc) of the rabbit antibodies and the Z domain moiety of the fusion protein. To obtain a signal useful for FACS, the enzymatic activity of the HRP conjugated to the rabbit  
10 antibodies was used through the addition of one ml of a signal amplification mixture containing fluorescein tyramide (Anton et al. J. Histochem. Cytochem. 46:771-777, 1998). Between each incubation step the beads were thoroughly washed, centrifuged for 3 min at  
15 2000 x g followed by resuspension in TNT buffer (0.1 M Tris-HCl pH 7.5, 0.15 M NaCl, 0.05% Tween 20) to remove non-specifically bound protein. TNB blocking buffer (0.1 M Tris-HCl pH 7.5, 0.15 M NaCl, 0.5% Blocking reagent from Tyramide Signal Amplification kit, NEN Life Science,  
20 Boston, MA, USA) was used during the incubation steps according to the manufacturers instructions.

After an incubation for five minutes at 25°C, and subsequent washing, the beads were resuspended in PBS for  
25 FACS analysis. This analysis showed that beads coated with the biotinylated BioM5 anti-FLAG antibody, incubated with the transcription/translation mixture of the FLAG-Z<sub>wt</sub> encoding PCR product could, subsequently incubated with the rabbit anti-DNP IgG-HRP conjugate and finally  
30 subjected to the signal amplification mixture containing fluorescein tyramide displayed significantly higher fluorescence signals in the FACS analysis than beads treated in the same way, but not containing the BioM5 anti-FLAG antibody (Figure 11).

35

This shows that fusion proteins, here exemplified by the fusion protein FLAG-Z<sub>wt</sub>, can be produced from a



corresponding PCR product by cell free transcription/  
translation containing a functional affinity fusion  
partner, here exemplified by the FLAG peptide, which is  
capable of resulting in a biospecific immobilization of  
5 the protein to beads containing a cognate affinity  
partner, here exemplified by the BioM5 anti-FLAG  
monoclonal antibody, and that such beads can be detected  
by FACS analysis using a suitable combination of  
detection reagents, here exemplified by a rabbit anti-DNP  
10 IgG-HRP conjugate and a signal amplification mixture  
containing fluorescein tyramide.

#### Example 6

Cell free transcription/translation of a bead-immobilized  
15 FLAG-Z<sub>wt</sub> encoding PCR product, biospecific immobilization  
of the gene product onto beads and analysis by  
fluorescence-activated cell sorting (FACS)

Biotinylated PCR fragments encoding a FLAG-Z<sub>wt</sub> fusion  
20 protein, obtained after PCR amplification using primers  
NOOL-12 and NOOL-13 on a plasmid pGEM-SD-K-FLAG-Z<sub>wt</sub>  
template were immobilized on streptavidin-coated beads  
(Bangs Laboratories) at a concentration of approximately  
10 ng/mg beads. The beads (50 mg) had previously been  
25 incubated with 5 µl of a solution containing 1.89 mg/ml  
of a biotinylated anti-FLAG peptide antibody (BioM5,  
Sigma). The beads containing both the biotinylated PCR  
products and the anti-FLAG peptide antibody were  
subjected to cell free transcription and translation  
30 using 25 ml of an S30 extract (Promega, Madison, WI,  
USA), supplemented with 200 units of T7 RNA polymerase  
(Epicentre, Madison, WI, USA) and 40 units of rRNasin  
(Promega, Madison, WI, USA). After incubation for one  
hour at 25°C, followed by repeated washing using TNT  
35 buffer (0.1 M Tris-HCl pH 7.5, 0.15 M NaCl, 0.05% Tween  
20), rabbit anti-DNP IgG antibodies conjugated to  
horse-radish peroxidase (HRP) (art. no. P0402, Dako,

- Denmark) were added to the beads and incubated overnight at 4°C (end-over-end mixing), followed by washing with TNT, to detect the translated and biospecifically immobilized FLAG-Z<sub>wt</sub> fusion protein gene product via the
- 5 biospecific interaction between the constant parts (Fc) of the rabbit antibodies and the Z domain moiety of the fusion protein (Nilsson et al., Protein engineering, 1: 107-113, 1987).
- 10 To obtain a signal useful for FACS, the enzymatic activity of the HRP conjugated to the rabbit antibodies was used through the addition of one ml of a signal amplification mixture containing fluorescein tyramide (Anton et al. J. Histochem. Cytochem. 46:771-777, 1998).
- 15 Between each incubation step the beads were thoroughly washed, centrifuged for 3 min at 2000 x g followed by resuspension in TNT buffer (0.1 M Tris-HCl pH 7.5, 0.15 M NaCl, 0.05% Tween 20) to remove non-specifically bound protein. TNB blocking buffer (0.1 M Tris-HCl pH 7.5,
- 20 0.15 M NaCl, 0.5% Blocking reagent from Tyramide Signal Amplification kit, NEN Life Science, USA) was used during the incubation steps according to the manufacturers instructions. As a negative control, streptavidin coated beads, containing immobilized BioM5 anti-FLAG antibodies,
- 25 and a PCR products obtained from PCR amplification using primers NOOL-12 and NOOL-13 on a plasmid pGEM-SD-K-FLAG-Z<sub>IgA</sub> template were included in the experiment.
- 30 The results from the FACS analysis shows that the beads containing the immobilized biotinylated PCR fragments encoding a FLAG-Z<sub>wt</sub> fusion protein, obtained after PCR amplification using primers NOOL-12 and NOOL-13 on a plasmid pGEM-SD-K-FLAG-Z<sub>wt</sub> template display a
- 35 significantly higher fluorescence intensity than the control beads containing immobilized PCR products

- 35 -

encoding a fusion protein not recognized by the reagent rabbit-HRP conjugate used for detection (Figure 12). This shows that fusion proteins, here exemplified by the fusion protein FLAG-Z<sub>wt</sub>, can be produced from a

5 corresponding, bead-immobilized, PCR product by cell free transcription/translation, containing a functional affinity fusion partner, here exemplified by the FLAG peptide, which is capable of resulting in a biospecific immobilization of the protein to beads containing a

10 cognate affinity partner, here exemplified by the BioM5 anti-FLAG monoclonal antibody, and that such beads can be detected by FACS analysis using a suitable combination of detection reagents, here exemplified by a rabbit anti-DNP IgG-HRP conjugate and a signal amplification mixture

15 containing fluorescein tyramide.

#### Example 7

#### Fluorescence-activated cell sorting (FACS)-based

20 enrichment of beads containing immobilized PCR products  
encoding a desired gene product

Biotinylated PCR fragments encoding FLAG-Z<sub>wt</sub> and FLAG-Z<sub>IgA</sub> fusion proteins, respectively, obtained after PCR amplification using primers NOOL-12 and NOOL-13 on

25 plasmids pGEM-SD-K-FLAG-Z<sub>wt</sub> and pGEM-SD-K-FLAG-Z<sub>IgA</sub> templates, respectively were separately immobilized on streptavidin-coated beads (Bangs Laboratories) to a level of approximately 10 ng/mg beads. The beads (50 mg) had previously been incubated with 5 µl of a solution

30 containing 1.89 mg/ml of a biotinylated anti-FLAG peptide antibody (BioM5, Sigma, Saint Louis, Mo, USA). Beads from the two pools were subsequently mixed at a ratio of 1:1 (equal amounts of beads of both sorts) and subjected to cell free transcription and translation using 25 ml of

35 an S30 extract (Promega, Madison, WI, USA), supplemented with 200 units of T7 RNA polymerase (Epicentre) and 40 units of rRNasin (Promega). After incubation for one

- 36 -

hour at 25°C, followed by repeated washing using TNT  
buffer (0.1 M Tris-HCl pH 7.5, 0.15 M NaCl, 0.05% Tween  
20), rabbit anti-DNP IgG antibodies conjugated to  
horse-radish peroxidase (HRP) (art. no. P0402, Dako,  
5 Denmark) were added to the beads and incubated for  
overnight at 4°C, followed by washing with TNT, to detect  
the translated and biospecifically immobilized FLAG-Z<sub>wt</sub>  
fusion protein gene product via the biospecific  
interaction between the constant parts (Fc) of the rabbit  
10 antibodies and the Z domain moiety of the fusion protein.

To obtain a signal useful for FACS, the enzymatic  
activity of the HRP conjugated to the rabbit antibodies  
was used through the addition of one ml of a signal  
amplification mixture containing fluorescein tyramide  
15 (Anton et al. J. Histochem. Cytochem. 46:771-777, 1998).

Between each incubation step the beads were thoroughly  
washed, centrifuged for 3 min at 2000 x g followed by  
resuspension in TNT buffer (0.1 M Tris-HCl pH 7.5, 0.15 M  
NaCl, 0.05% Tween 20) to remove non-specifically bound  
20 protein. TNB blocking buffer (0.1 M Tris-HCl pH 7.5,  
0.15 M NaCl, 0.5% Blocking reagent from Tyramide Signal  
Amplification kit, NEN Life Science, Boston, MA, USA) was  
used during the incubation steps according to the  
manufacturers instructions. Using FACS, a bead pool  
25 originally obtained by the mixing at the 1:1 bead ratio  
was subsequently subjected to enrichment experiment based  
on fluorescence intensity. In this procedure the  
settings in the FACS instrument were adjusted for  
preparative isolation of single beads (singlets) having a  
30 relative fluorescence intensity above 50. With this  
setting, the mixture was subjected to sorting and tubes  
with approximately 4500 sorted beads were collected.

To analyze if beads carrying the PCR products encoding  
35 the FLAG-Z<sub>wt</sub> fusion protein, which should be specifically  
labeled by the labeling procedure involving the rabbit  
IgG-HRP conjugate, were enriched relative to beads

- 37 -

carrying the PCR products and FLAG-Z<sub>IgA</sub> fusion proteins not being recognized by the rabbit IgG-HRP conjugate, the difference in DNA sequence between the two PCR products was employed.

5

The FLAG-Z<sub>wt</sub> fusion protein-encoding PCR products contain a recognition sequence for the enzyme Mlu I, not present in the PCR products encoding the FLAG-Z<sub>IgA</sub> fusion protein. This allowed for a discrimination between the two PCR products through an analysis of the susceptibility for Mlu I digestion (Figure 13A). Samples of beads from before and after sorting were therefore subjected to PCR amplification using primers NOOL-12 and NOOL-13, which anneals at sites in the immobilized PCR products flanking the regions which differs between the two PCR product species, and therefore could be use for the simultaneous amplification of both PCR product species. Subsequent incubation of the resulting new PCR products with the restriction enzyme Mlu I could therefore be used to investigate the relative ratios between the two species in samples from before and after sorting, by analysis of DNA fragment sizes and band intensities after agarose gel electrophoresis followed by ethidium bromide staining.

25

A PCR amplification of the nucleic acids present on approximately 10000 beads from the 1:1 mixture (sample from before sorting) followed by a digestion with Mlu I and analysis by gel electrophoresis shows, as expected, upon a mixture of Mlu I-susceptible and Mlu I-resistant PCR products (Figure 13B, lane 6).

30

When approximately 400 beads collected during the FACS enrichment was subjected to the same analysis, the intensity ratio between the upper band (443 bp, uncleaved) and lower double band (two cleavage products, 239/204 bp, unresolved) had shifted towards the smaller

35

(lower) bands (Figure 13B, lane 8). Using a Gel Doc 2000 gel scanning instrument and Quantity One vers. 4.1 software (Biorad, Hercules, CA, USA), this shift in relative intensities were recorded resulting in the overlay plot shown in figure 14. From this analysis it can be clearly seen that a shift of the relative intensity towards the lower molecular weight cleavage products had occurred. This shows that beads containing Mlu I-susceptible PCR product encoding the FLAG-Z<sub>wt</sub> fusion protein, had been enriched during the experiment, relative to beads containing the Mlu I-resistant FLAG-Z<sub>IgA</sub> fusion protein encoding PCR product.

Taken together, this example shows that fusion proteins, here exemplified by the fusion protein FLAG-Z<sub>wt</sub>, can be produced from a corresponding, bead-immobilized, PCR product by cell free transcription/translation, containing a functional affinity fusion partner, here exemplified by the FLAG peptide, which is capable of resulting in a biospecific immobilization of the protein to beads containing a cognate affinity partner, here exemplified by the BioM5 anti-FLAG monoclonal antibody, and that such beads can be enriched when mixed and co-processed with irrelevant beads, containing PCR products encoding a different gene product, by FACS-based enrichment using a suitable combination of detection reagents, here exemplified by a rabbit anti-DNP IgG-HRP conjugate and a signal amplification mixture containing fluorescein tyramide.